arXiv:1212.3118v2 [hep-ph] 20 Dec 2012

Magnetic test of chiral dynamics in QCD

Yu.A.Simonov,
Institute of Theoretical and Experimental Physics
117118, Moscow, B.Cheremushkinskaya 25, Russia

Abstract

Magnetic fields in the range $0 \leq eB \lesssim 1$ GeV^2 effectively probe internal quark structure of chiral mesons and test basic parameters of chiral theory, such as $\langle \bar{q}q \rangle, f_\pi$. We extend previously formulated chiral theory, including quark degrees of freedom, to the case of strong magnetic fields and show that the quark condensate $|\langle \bar{q}q \rangle|_{u,d}$ grows quadratically with eB for $eB \lesssim 0.2$ GeV^2 and linearly for higher field values. These results agree quantitatively with recent lattice data and differ from χPT predictions.

1 Introduction

Strong magnetic fields (MF) are expected to be present in the early universe [1, 2, 3], in neutron stars (magnetars) [4, 5], and also in non-central heavy-ion collisions [6]. It is very interesting how MF can modify strong interactions, deduced from QCD. In particular, it was recently shown [7], how MF influence asymptotic freedom behavior of the QCD strong coupling constant and gluon exchange interaction. The interplay of confinement and MF in neutral mesons was studied in [8], showing a dramatic change of masses with growing MF.

Of a particular interest is the influence of MF on the chiral symmetry breaking (CSB) dynamics, and in a more general setting, on symmetry breaking pattern in field theory.

It was emphasized in [9], that MF has the tendency to strengthen the chiral condensate, the phenomenon called there the magnetic catalysis, see [10] for a review.

The behavior of chiral condensate in MF was studied in the chiral perturbation theory (χPT) [11, 12, 14], in Polyakov-Nambu-Jona-Lasinio (PNJL) [15] and in other models [16]. On the lattice the corresponding analysis was done for quenched QCD in [17, 18], in $n_f = 2\ QCD$ in [19, 20] and in $n_f = 4$ SU(2) theory in [21]. In all cases both in models and on the lattice the chiral condensate grows with MF, however in different way.

Recently a comprehensive analysis of this problem on the lattice with physical quark masses was performed in [22] and has shown a quadratic growth of condensate up to $eB \sim 0.2$ GeV^2 and approximately linear behavior above this value.

This linear behavior contradicts the (χPT) results, see e.g. [11, 14], where linear growth was found with the slope almost twice as small as in [22], and also contradicts PNJL quadratic growth at large eB [15].

As it was mentioned in [13], the reasonable region of χPT application is for $eB \lesssim m_\pi^2$, where the quark condensate grows quadratically, in qualitative agreement with lattice data however the linear behavior up to 1 GeV^2 is outside of χPT reliable region.

It is the purpose of the present paper to study quark condensate in MF starting from the basic QCD equations, derived earlier without MF in [23, 24, 25, 26]. It was shown there, that GMOR relations and expressions for $\langle \bar{q}q \rangle$ and f_π can be derived from the basic QCD quantities: string tension, α_s and current quark masses in good agreement with experiment and lattice data.

Recently [27] these results were extended to account for growing current quark masses m_q and in particular the dependence of $\langle \bar{q}q \rangle$ on m_q was established to be in agreement with lattice data [28].

In the present paper we follow the same line of the formalism of [23, 24, 25, 26], but now in MF and find the behavior of $\langle \bar{q}q \rangle$ for u and d quarks and their average with growing MF.

As a result we observe in the resulting dependence of the average $\langle \bar{q}q \rangle$ and $\langle \bar{u}u \rangle$, $\langle \bar{d}d \rangle$, the same features and good quantitative agreement with the lattice data obtained in [22]. The physical reason for this dependence of $\langle \bar{q}q \rangle$ on eB is clarified below in the paper.

The paper is organized as follows. In the next section a general derivation of GMOR relation and expressions for $\langle \bar{q}q \rangle$ and f_π are given, in section 3 the MF dependence of basic terms is established, in section 4 results are discusse and prospectives are given.

2 Derivation of the effective chiral lagrangian in MF

We follow here [24] to write first the effective lagrangian of the light quark in the confining field of the antiquark in MF, starting with the standard QCD partition function in Euclidean space-time

$$Z = \int DAD\psi D\psi^+ \exp[L_0 + L_1 + L_{int}] \tag{1}$$

where

$$L_0 = -\frac{1}{4}\int d^4x (F^a_{\mu\nu})^2, \tag{2}$$

$$L_1 = -i\int {}^f\psi^+(x)(\hat{D}+m_f)^f\psi(x)d^4x \tag{3}$$

$$L_{int} = \int {}^f\psi^+(x)g\hat{A}(x)^f\psi(x)d^4x \tag{4}$$

and

$$\hat{D} = \gamma_\mu(\partial_\mu - ieA^{(e)}_\mu(x)), \quad A^{(e)}_\mu(x) = \frac{1}{2}[\mathbf{x}\times\mathbf{B}]. \tag{5}$$

Averaging Z over vacuum gluonic field and keeping only lowest (bilocal) correlators of color fields $D_{\mu\nu,\lambda\sigma}(x,y) = \frac{1}{N_c} tr\langle F_{\mu\nu}(x)\Phi(x,y)F_{\lambda\sigma}(y)\Phi(y,x)\rangle$, one finds as in [24],

$$\langle Z\rangle_A = \int D\psi D\psi^+ \exp(L_1 + L^{(4)}_{eff}), \tag{6}$$

where

$$L^{(4)}_{eff} = \frac{1}{2N_c}\int d^4x d^4y\ {}^f\psi^+_{a\alpha}(x){}^f\psi_{b\beta}(x){}^g\psi_{b\gamma}(y){}^g\psi_{a\delta}(y)J_{\alpha\beta,\gamma\delta}(x,y) \tag{7}$$

and

$$J_{\alpha\beta,\gamma\delta}(x,y) = (\gamma_\mu)_{\alpha\beta}(\gamma_\nu)_{\gamma\delta}J_{\mu\nu}(x,y) \tag{8}$$

$$J_{\mu\nu}(x,y) = g^2\int_C^x du_\alpha \int_C^y dv_\beta D_{\alpha\mu,\beta\nu}(u,v) \tag{9}$$

Here indices f,g refer to flavor, a,b to color and α,β,μ,ν to Lorentz indices. Eq. (9) implies that some contour gauge is used for simplicity, but the final result is gauge invariant and the most important property of $J_{\mu\nu}(x,y)$ is that it is proportional to the distance of the average point $\left(\frac{x+y}{2}\right)$

to the contour C (linear confinement), and the effective distance between x and y (nonlocality) is of the order of the vacuum correlation length $\lambda \approx 0.1$ fm.

In the large N_c limit the four-quark expression in $L^{(4)}_{eff}$ can be replaced by the quadratic one, using the limit

$$ {}^f\psi_{b\beta}(x){}^g\psi_{b\gamma}(y) \to \delta_{fg} N_c\ {}^f S_{\beta\gamma}(x,y), \tag{10}$$

where ${}^f S_{\beta\gamma}(x,y)$ is the quark propagator.

As a result one obtains the form

$$L^{(4)}_{eff} \to -i \int d^4x d^4y\ {}^f\psi^+_{a\alpha}(x)^{(fg)} M_{\alpha\beta}(x,y)\ {}^g\psi_{a\beta}(y), \tag{11}$$

$$ {}^{fg}M_{\alpha\delta}(x,y) = -iJ_{\mu\nu}(x,y)(\gamma_\mu\ {}^{fg}S(x,y)\gamma_\nu)_{\alpha\delta}, \tag{12}$$

and the quark propagator satisfies the equation

$$(-i\hat{D} - im_f)\ {}^f S(x,y) - i\int\ {}^{(fg)}M(x,z)\ {}^g S(z,y) = \delta^{(4)}(x,y). \tag{13}$$

It is convenient to use the following parametrization of ${}^f M(x,y)$ in terms of scalar functions, flavor singlet $M_s(x,y)$ and flavor triplet $\phi_a(x,y), a = 1,2,3,$

$$M^{(fg)}_{\alpha\beta} x,y) = M_s(x,y)\exp(i\gamma_5 t^a \phi_a(x,y))^{(fg)}_{\alpha\beta} \equiv M_s(x,y)\hat{U}^{(fg)}_{\alpha\beta}(x,y) \tag{14}$$

As a result the effective Lagrangian assumes the form

$$L_\phi = \int d^4x d^4y \left\{ {}^f\psi^+_{a\alpha}(x)[(i\hat{D} + im_f)_{\alpha\beta}\delta^{(4)}(x-y)\delta_{fg} + iM_s\hat{U}^{(fg)}_{\alpha\beta}(x,y)]\ {}^g\psi_{a\beta}(y)\right\}, \tag{15}$$

and the partition function can be written as

$$\langle Z\rangle_A = \int D\psi D\psi^+ DM_s D\phi_a \exp L_\phi. \tag{16}$$

Integrating over $D\psi D\psi^+$ one obtains the effective chiral Lagrangian (ECL) L_{ECL},

$$\langle Z\rangle_A = \int DM_s D\phi_a \exp L_{ECL}, \tag{17}$$

where

$$L_{ECL} = N_c trlog[(i\hat{D} + im_f)\hat{1} + iM_S\hat{U}]. \tag{18}$$

Finally,the ECL at the stationary point in the integral (17) is defined by conditions $\frac{\delta L_{ECL}}{SM_S} = \frac{\delta L_{ECL}}{\delta\phi_a} = 0$,which yields

$$iM_s^{(0)}(x,y) = (\gamma_\mu S\gamma_\nu)J_{\mu\nu}(x,y), \quad \phi_a^{(0)} = 0, \tag{19}$$

and

$$S = S_\phi(\phi_a = 0), \quad S_\phi = -[(i\hat{D} + im_f)\hat{1} + iM_s\hat{U}].$$

Insertion of (19) in (18) yields the effective action for pseudoscalar fields ϕ_a

$$L_{ECL} \to -W(\phi) = N_c tr\log[(i\hat{D} + im_f)\hat{1} + iM_s\hat{U}]. \tag{20}$$

Our final step here is the local limit of $J_{\mu\nu}(x,y)$ and $M_s^{(0)}(x,y)$ proved in [23], which yields

$$\phi_a(x,y) \to \phi_a(x), \quad M_s^{(0)}(x,y) \to M_s^{(0})(x)\delta^{(4)}(x-y) \tag{21}$$

Expanding $W(\phi)$ in powers of ϕ_a and keeping quadratic terms, one has

$$W^{(2)}(\phi) = \frac{1}{2}\int \frac{d^4kd^4k'}{(2\pi)^4(2\pi)^4}\phi_a(k)N(k,k')\phi_a(k'), \tag{22}$$

where

$$\hat{N}(k,k') = \frac{N_c}{2}\int dx e^{i(k+k')x} tr(\Lambda M_s)_{xx} +$$

$$\frac{N_c}{2}\int d^4(x-y)d^4\left(\frac{x+y}{2}\right)\exp\left[i\frac{(k+k')}{2}(x+y) + \frac{i}{2}(k-k')(y-x)\right]\times$$

$$\times\, tr[\Lambda(x,y)M_s(y)\bar{\Lambda}(y,x)M_s(x)], \tag{23}$$

with the definitions

$$\Lambda = (\hat{D} + m + M_s)^{-1}, \quad \bar{\Lambda} = (\hat{D} - m - M_s)^{-1}. \tag{24}$$

Using translation invariance of traces in (23) one can rewrite it as

$$W^{(2)}(\phi) = \frac{N_c}{2}\int \phi_a(k)\phi_a(-k)\bar{N}(k)\frac{d^4k}{(2\pi)^4}, \tag{25}$$



Figure 1: Quadratic terms in the pionic action (18)

where

$$\bar{N}(k) = \frac{1}{2} tr\{(\Lambda M_s)_0 + \int d^4 z e^{ikz} \Lambda(0,z) M_s(0) \bar{\Lambda}(z,0) M_s(0)\} = \frac{1}{2} tr(\Lambda M). \tag{26}$$

The graphical representation of $W^{(2)}(\phi)$ is given in Fig.1 and the picture, which helps to understand the physical meaning of $M(x,y)$ is given in Fig.2. As a result the standard chiral condensate in the Minkowskian space-time can be written as

$$\langle \bar{\psi}\psi \rangle_M = i\langle \psi\psi^+ \rangle_E = -N_c tr\Lambda = \langle \bar{u}u + \bar{d}d \rangle_M. \tag{27}$$

This can be rewritten as

$$tr\Lambda = -(M(0)+m) G^{(0)}(k=0), \tag{28}$$

where $m = \frac{m_u+m_d}{2}$, and $M(0)$ is expressed via the vacuum correlation length λ [24]

$$M(0) = \frac{2}{\sqrt{\pi}} \sigma\lambda(1 + O(\sigma\lambda^2) \tag{29}$$

Here λ is the vacuum correlator length, defined from the mass in the field correlator [29], 0.1 fm $\lesssim \lambda \lesssim$ 0.2 fm. Therefore 100 MeV $\leq M(0) \leq$ 200 MeV.

Writing the Green's function $G^{(0)}(k)$ as a spectral decomposition, one obtains

$$G^{(0)}(k) = -\sum_{n=0} \frac{c_n^2}{k^2 + m_n^2}. \tag{30}$$

In a similar way, as shown in [24] and [30], the $q\bar{q}$ Green's function $G^{(MM)}(k)$ with $M(0)$ in both vertices is connected to decay constant f_π,



Figure 2: The flux-tube operator $M(x,y)$ in the Wilson loop

namely

$$G^{(MM)}(k)=-M^2(0)\sum_n\frac{c_n^2}{k^2+m_n^2},\quad G^{(MM)}(k)-G^{(MM)}(0)=\frac{k^2f_\pi^2}{2n_c}+O(k^4). \tag{31}$$

One can express c_n as $c_n=\sqrt{\frac{m_n}{2}}\psi_n(0)$ and finally one obtains chiral relations for $B=0$, see [24] for details of derivation

$$-\langle\bar{q}q\rangle_i=N_c(M(0)+m_i)\sum_{n=0}^{N}\frac{\psi_n^2(0)}{m_n}e^{-m_n\lambda},\quad i=u,d,s \tag{32}$$

$$f_\pi^2=N_cM^2(0)\sum_{n=0}^{N}\frac{\psi_n^2(0)}{m_n}e^{-m_n\lambda}(1+m_n\lambda). \tag{33}$$

Note, that the GMOR relation [31] is modified,

$$m_\pi^2f_\pi^2=\frac{\bar{m}M(0)}{M(0)+\bar{m}}|\langle\bar{u}u\rangle+\langle\bar{d}d\rangle|,\quad \bar{m}=\frac{m_n+m_d}{2}, \tag{34}$$

however for $\bar{m}\to 0$, (34) coincides with the original form.

Eqs. (32), (33) express quark condensate and f_π in terms of $M(0)$ and the reference spectrum in the PS channel, i.e. the spectrum, where chiral symmetry is not taken into account, but only confinement term $M_s(x)$ is present, the corresponding masses m_n, calculated in [24] are $m_0=0.4$ GeV, $m_1=1.35$ GeV, $m_2=1.85$ GeV. Taking $\lambda=0.1$ fm and $M(0)=0.15$ GeV, one obtains (cf appendix of the second paper in [24])

$$-\frac{\langle\bar{q}q\rangle}{n_f}=(217\text{MeV})^3\frac{M(0)+\bar{m}}{(150\text{MeV})},\quad f_\pi=96\text{GeV}\frac{M(0)+\bar{m}}{(150\text{MeV})}. \tag{35}$$

One can see, that our values are in the correct ballpark, close to average lattice and phenomenological values.

Now we turn to the case of nonzero MF. Eq. (32) in case of MF can be rewritten for each flavor separately,

$$|\langle\bar{q}q\rangle_i|=N_c(M(0)+m_i)\sum_{n=0}^{\infty}\left(\frac{\frac{1}{2}|\psi^{(+-)}_{n,i}(0)|^2}{m^{(+-)}_{n,i}}+\frac{\frac{1}{2}|\psi^{(-+)}_{n,i}(0)|^2}{m^{(-+)}_{n,i}}\right) \tag{36}$$

where $i=u,d,s$ and the subscripts $(+-)$ and $(-+)$ refer to the quark and antiquark spin projections on the MF $\mathbf{B}$, and the coefficients $\frac{1}{2}$ are due to γ_5 in the vertices of the Green's function $G^{(0)}(k)$, $\langle\gamma_5|\to\frac{\langle+-|-\langle-+|}{\sqrt{2}}$.

The main problem reduces to the calculation of the spectrum of eigenvalues and eigenfunctions ψ_n, m_n, which are to be found from the Hamiltonian H containing MF $\mathbf{B}$, and derived from the path integral Hamiltonian [8]

$$H_0=\sum_{i=1}^{2}\left(\frac{(\mathbf{p}_i-e_i\mathbf{A}^{(e)})^2}{2\omega_i}+\frac{m_i^2+\omega_i^2}{2\omega_i}-\frac{e_i\boldsymbol{\sigma}_i\mathbf{B}}{2\omega_i}\right) \tag{37}$$

which can be rewritten in the form $(i=u,d,s)$

$$H_i=\frac{1}{2\tilde{\omega}}\left(-\frac{d^2}{d\boldsymbol{\eta}^2}+\left(\frac{e_i(\boldsymbol{\eta}\times\mathbf{B})}{2}\right)^2\right)+V_{\text{conf}}(\eta)-\sum_{k=1,2}\frac{e_i\boldsymbol{\sigma}_k\mathbf{B}}{2\omega_k}, \tag{38}$$

where $\tilde{\omega}=\frac{\omega_1\omega_2}{\omega_1+\omega_2}$.

The eigenvalues $\tilde{m}_{n,i}$ of H_i depend on ω_1,ω_2, and the final eigenvalues $m_{n,i}$ entering in (3) are obtained as stationary points in variation over ω_1,ω_2

$$\left.\frac{\partial\tilde{m}_{n,i}}{\partial\omega_{1,2}}\right|_{\omega^{(0)}_{1,2}}=0,\quad \text{e.g. } \tilde{m}^{(+-)}_{n,i}(\omega^{(0)}_1,\omega^{(0)}_2)\equiv m^{(+-)}_{n,i}. \tag{39}$$

To avoid purely numerical calculations, one can simplify the hamiltonian (38), replacing linear confinement $V_{\text{conf}}(\eta)$ in (38) by a suitable quadratic form, with coefficient found from the stationary point condition

$$V^{(lin)}_{conf}=\sigma\eta\to V^{(\text{quadr})}_{\text{conf}}=\frac{\sigma}{2}\left(\eta^2\gamma+\frac{1}{\gamma}\right),\quad \left.\frac{\partial\tilde{m}_{n,i}}{\partial\gamma}\right|_{\gamma=\gamma_0}=0 \tag{40}$$

To check the accuracy of this replacement, one can compare the lowest mass eigenvalues and $\psi_n(0)$ first without MF. In particular, $m_0^{(\rm quadr)} = 2\sqrt{3\sigma}$, while $m_0^{\rm lin} = 4\sqrt{\sigma}\left(\frac{2.338}{3}\right)^{3/4}$, and these two figures differ by 4.5%.

For $\psi_n(0)$ the corresponding results are

$$|\psi_n(0)|^2_{\rm lin} = \frac{0.82\sigma^{3/2}}{4\pi} = 0.065\sigma^{3/2};$$

$$|\psi_0(0)|^2_{\rm quadr} = \frac{\sigma^{3/2}(\frac{c_0}{4})^{3/4}}{\pi^{3/2}} = 0.065\sigma^{3/2}(c_0 \equiv 1) \tag{41}$$

This coincidence of $|\psi_0(0)|^2$ will be of special importance in what follows, since the main effect of MF, as will be seen, is the increase of $|\psi_n(0)|^2$ in (36) due to MF. It is clear, that with growing B the size of meson is decreasing, and the difference between $V^{(\rm lin)}_{\rm conf}$ and $V^{(\rm quadr)}_{\rm conf}$ will be even more suppressed, since these both interactions vanish simultaneously, $V^{(\rm lin)}_{\rm conf}(r \to 0) = V^{(\rm quadr)}_{\rm conf}(r \to 0) = 0$.

For us it will be most important how $|\psi_n(0)|^2$ depends on MF, and especially, how MF enters in the expansion of $|\psi_n(0)|^2$ in powers of B. To this end one can see in the Hamiltonian (38), that MF enters via the term $V_B \equiv \frac{\boldsymbol{\eta}_\perp^2}{2\tilde{\omega}}\left(\frac{e_i\mathbf{B}}{2}\right)^2$. It is clear, that in the perturbative series expansion the MF enters as $(e_iB)^2$

$$\psi_n(0)_B = \psi_n(0)_0 + O((e_iB)^2) + ... \tag{42}$$

As a result of (40) one can immediately write the analytic expressions for $\psi_n(0)$ and m_n of the following form, e.g.

$$|\psi_n(0)|^2 = \frac{1}{\pi^{3/2}r_\perp^2 r_3}, \quad r_\perp^2 = 2\left((e_qB\ )^2 + \sigma^2 c\right)^{-1/2}, \quad r_3 = \left(\frac{\sigma^2 c}{4}\right)^{-1/4} \tag{43}$$

where $c = \frac{4\tilde{\omega}}{\gamma\sigma}$, and

$$m_{n_\perp,n_3} = \varepsilon_{n_\perp,n_3} + \frac{m_1^2 + \omega_1^2 - e_q\mathbf{B}\boldsymbol{\sigma}_1}{2\omega_1} + \frac{m_2^2 + \omega_2^2 + e_q\mathbf{B}\boldsymbol{\sigma}_2}{2\omega_2} \tag{44}$$

where

$$\varepsilon_{n_\perp,n_3} = \frac{1}{2\tilde{\omega}}[\sqrt{e_q^2B^2 + \sigma^2 c}(2n_\perp + 1) + \sqrt{\sigma^2 c}(n_3 + \frac{1}{2})] + \frac{\gamma\sigma}{2} \tag{45}$$

Expressions (36),(43), (44), (45) contain all information necessary to computer quark condensate for varying MF.

3 The MF dependence of quark condensates

We are interested in eigenfunctions and eigenvalues of $(+-)$ and $(-+)$ states both for u and d quarks. For the $(+-)$ state, one can see in (45), (44) that at large B, $m^{(+-)}_{0,n_3}$ tends to a constant limit, together with $\omega_1 = \omega_2$. In this case the parameter c in (43) and (45) which can be expressed in general as

$$c_{+-} = \frac{4\tilde{\omega}}{\gamma\sigma} = \left(\frac{a}{2}\right)^{4/3} 4/\beta;\quad \gamma(B) = \beta(B)(\sigma\tilde{\omega})^{-1/3},\quad \omega(B) = a(B)\sqrt{\sigma} \tag{46}$$

and $\beta(B), a(B)$ are changing in finite limits and one obtains that $c_{+-}(B) \approx 1$, for all B, and m_{+-} tends to a constant limit for $B \to \infty$. Hence one can write for lowest levels $n_3 = 0, 1, 2, \ldots$

$$|\psi^{(+-)}_{n_\perp=0,n_3}(0)|^2 \cong \frac{\sqrt{\sigma}\sqrt{e_q^2 B^2 + \sigma^2}}{(2\pi)^{3/2}}. \tag{47}$$

Note, that in the limit $B \to 0$ this expression for $|\psi(0)|^2$ yields equal values for $n_3 = 0, 1, 2$ as it should be for pure linear confining interaction.

For the $(-+)$ case the situation is different, and at large B the stationary point value $\omega_0^{(-+)} \approx \sqrt{2|e_q|B^+ + \frac{\sigma}{4}}$, and the parameter c_{-+} is increasing with B:

$$c_{-+}(B) = \left(1 + \frac{8 e_q B}{\sigma}\right)^{2/3}. \tag{48}$$

As a result, the $|\psi_n^{(-+)}(0)|^2$ can be written as

$$|\psi_{n_3}^{(-+)}(0)|^2 = (\sigma^2 c_{-+})^{3/4} \sqrt{1 + \left(\frac{e_q B}{\sigma}\right)^2 \frac{1}{c_{-+}}}. \tag{49}$$

Moreover, $m^{(-+)}_{n_3} \approx 2\sqrt{2|e_q B| + \frac{\sigma}{4}}$, and $\frac{(\sigma^2 c_{-+})^{3/4}}{m_0^{(-+)}(B)} = \sigma$ at large B.

Combining (45) and (49) we therefore can write

$$|\langle \bar{q}q\rangle_i(B)| = |\langle \bar{q}q\rangle_i(0)| \frac{1}{2} \left\{ \sqrt{1 + \left(\frac{e_q B}{\sigma}\right)^2} + \sqrt{1 + \left(\frac{e_q B}{\sigma}\right)^2 \frac{1}{c_{-+}}} \right\} \tag{50}$$

where c_{-+} is given in (48).

Table 1: Values of $\Delta \sum_i(B), i = u, d$ given by (50) in comparison with lattice data from [22]

eB (GeV^2)	0	0.2	0.4	0.6	0.8	1
$\Delta \sum_u$ this paper	0	0.156	0.48	0.865	1.273	1.65
$\Delta \sum_u$ [21] lattice	0	0.185	0.51	0.86	1.235	1.60
$\Delta \sum_d$ this paper	0	0.048	0.158	0.308	0.48	0.67
$\Delta \sum_d$ [21] lattice	0	0.095	0.23	0.40	0.57	0.73

We shall be using notations of [22] for the increment of quark condensate as a function of MF (denoted $\tau_i(B)$ in [9])

$$\Delta \sum_i(B) = \frac{|\langle \bar{q}q\rangle_i(B)| - |\langle \bar{q}q\rangle_i(0)|}{|\langle \bar{q}q\rangle_i(0)|}. \tag{51}$$

The resulting values of $\Delta \sum_i(B)$ from (50), (51) are compared in Table 1 with the corresponding lattice calculations in [22].

One can see a reasonable agreement between our theory and lattice data. One can simplify the B dependence of Eq. (50), writing

$$|\langle \bar{q}q\rangle_i(B)| = |\langle \bar{q}q\rangle_i(0)|\sqrt{1 + \left(\frac{e_i B}{M_i^2}\right)^2}. \tag{52}$$

The form (52) also satisfactorily describes data of [22] and [20], taking M_i as a fitting parameter. In the case of [20]the fitted values of M_i are approximately

$$M_u^2 \approx 0.35\text{GeV}^2, \quad M_d^2 \approx 0.27\text{GeV}^2, \tag{53}$$

i.e. are 2 and 1.5 times larger, than σ (the corresponding $c = c_{+-} = c_{-+} \cong 4$ and 4.25 for u and d cases), however the agreement with lattice data is worse. Qualitatively the same situation (with even larger M_i^2)takes place in comparison with [17, 18].

As a whole the behavior (50),(52) correctly reproduces the main qualitative features of the quark condensate as a function of eB: the quadratic behavior proportional to $(e_iB)^2$ at small B and linear behavior $\sim |e_iB|$ at large B. It differs from the results of others approaches. In particular, χPT [11] predicts linear behavior in eB with the slope much smaller, than in lattice data [22].

As it was mentioned in [20], for $\Delta\sum_u$ and $\Delta\sum_d$ there is a simple relation

$$\Delta\sum_u\left(\frac{B}{2}\right) = \Delta\sum_d(B), \tag{54}$$

which is satisfied in lattice data, and, of course, is trivially satisfied in our definitions.

Writing for large $B \gg \sigma$, that $\Delta\sum_i(B) \cong a_iB$, one immediately obtains from (54), that $a_u = 2a_d$. This relation is approximately satisfied in lattice data [20, 22], and in our expression (50), (52).

4 Discussion and conclusions

We have used our formalism for chiral dynamics, presented in [23, 24, 25, 26, 27], which is derived not from purely symmetry considerations, but from the QCD quark dynamics, where chiral symmetry appears approximately in the small m_q limit of the effective QCD Lagrangian. In this way all basic degrees of freedom are connected to the confined quarks, and it is finally the confinement, which dictates properties of fundamental chiral quantities $\langle\bar{q}q\rangle$, f_π etc., and gives them numerical values, expressed via σ. This is in contrast with standard chiral ideas, where chiral constants are postulated from the beginning and chiral symmetry breaking is not connected to the basic QCD quark dynamics.

Actually our reasoning for the calculation of quark condensate is very simple. After one derives scalar confining interaction $M_s(x)$, which acts on each quark or antiquark, one can write quark condensate $\langle\bar{q}q\rangle = -N_c tr\Lambda$ as

$$tr\Lambda = \left\langle tr\frac{1}{M_s+m_q+\hat{\partial}}\right\rangle = \left\langle tr\left[\frac{1}{M_s+m_q+\hat{\partial}}(M_s+m_q-\hat{\partial})\frac{1}{(M_s+m_q-\hat{\partial})}\right]\right\rangle =$$

$$= (M_s(0)+m_q)\left\langle tr\left[\gamma_5\frac{1}{M_s+m_q+\hat{\partial}}\gamma_5\frac{1}{M_s+m_q+\hat{\partial}}\right]\right\rangle, \tag{55}$$

since the term with $\hat{\partial}$ is odd and vanishes.

The last quantity on the r.h.s. of (55) is the $q\bar{q}$ Green's function, proportional to $\psi_n^2(0)$ for each n state. The only difference in presence of MF is the replacement $\hat{\partial} \to (\hat{\partial} - ie\hat{A}^{(e)})$, which immediately gives the proportionality $\langle \bar{q}q \rangle \sim \psi^2(r=0; eB)$. This latter quantity is linearly rising with $|eB|$, since MF is "focussing" quark-antiquark system at small distances.

External magnetic field is here crucial for our understanding of chiral dynamics, and using MF one may decide, what is the role of quark dynamics in the chiral phenomena. In this respect the comparison of the behavior of quark condensate (or $\Delta \sum(B)$) as a function of B in different models and lattice data is showing the following:

1. Standard chiral theory at large $B \sim \sigma$ gives linear behavior (qualitatively correct) but with the wrong slope.

2. Our approach, Eq. (36) predicts quadratic behavior at small B $B \lesssim \sigma$ and linear at larger B with slopes different for u and d quarks, both in agreement with existing lattice data.

3. The PNJL model as shown in [22], is also in disagreement with lattice data at larger eB.

Physically, it is clear, that MF, acting on quark charges, immediately discloses the internal quark structure of PS mesons, while the effective chiral Lagrangian, as in [11, 12] describes only internal multipionic degrees of freedom. The latter can be important only for $B \ll 1/r_0^2$, where $r_0 \approx 0.6$ fm is the pionic radius, i.e. for $B \ll 0.1$ GeV^2, while for large B the standard chiral picture is irrelevant, as it is confirmed by lattice data.

These considerations suggest the idea, that the true chiral dynamics can be derived e.g. from the effective Lagrangian (18), and should finally display coexisting quark and chiral symmetric degrees of freedom, demonstrating how the latter disappear (suppressed) for growing quark masses. How it happens with the pionic spectra, was explained in [27], demonstrating the unifying spectrum based on chiral and quark degrees of freedom at the same time.

The author is grateful for discussions to N.O.Agasian, M.A.Andreichikov and B.O.Kerbikov.

References

[1] T.Vachaspati, Phys. Lett. **B265**, 258(1991).

[2] K.Enquist and P.Olesen, Phys. Lett. B **319**, 178 (1993).

[3] D.Grasso and H.R.Rubinstein, Phys. Rept. **348**, 163 (2001).

[4] R.C.Duncan and C.Thompson, Astro. Phys. J. **392**, L9 (1992).

[5] J.M.Lattimer and M.Prakash, Phys. Rept. **442**, 109 (2007); A.K. Harding and Dong Lai, Rep. Progr. Phys. **69**, 1831 (2006).

[6] D.E.Kharzeev, L.D.McLerran and H.J.Warringa, Nucl. Phys. A **803**, 227 (2008); V.S.Skokov, A.Illarionov and V.Toneev, Int. J. Mod. Phys. A **24**, 5925 (2009).

[7] M.A.Andreichikov, V.D.Orlovsky and Yu.A.Simonov, arXiv: 1211.6568 [hep-ph].

[8] M.A.Andreichikov, B.O.Kerbikov and Yu.A.Simonov, arXiv:1210.0227, [hep-ph].

[9] V.Gusynin, V.Miransky and I.Shovkovy, Phys. Rev. Lett. **73**, 3499 (1994).

[10] I.A.Shovkovy, arXiv: 1207.5081.

[11] I.Shushpanov and A.V.Smilga, Phys. Lett. B **402**, 351 (1997).

[12] N.O.Agasian and I.Shushpanov, Phys. Lett. B**472**, 143 (2000); N.O.Agasian, Phys. Lett. B **488**, 39 (2000); J.O.Andersen, arXiv;1205.6978, JHEP 1210 (2012) 005.

[13] N.O.Agasian, Phys. Atom. Nucl. **64**, 554 (2001).

[14] T.D.Cohen, D.A.Mc Gady, and E.S.Werbos, Phys. Rev. C **76**, 055201 (2007); J.O.Andersen, arXiv:1202.2051; 1205.6978.

[15] R.Gatto and M.Ruggieri, Phys. Rev. D **83**, 034016 92011).

[16] N.O.Agasian and S.M.Fedorov, Phys. Lett. B **663**, 445 (2008); E.S.Fraga and A.J.Mizher, Nucl. Phys. A **820**, 034016 (2011); E.S.Fraga and L.F.Palhares, arXiv:1201.5881.

[17] P.Buividovich, M.Chernodub, E.Luschevskaya and M.Polikarpov, Phys. Lett. B **682**, 484 (1010); arXiv:0812.1740.

[18] V.Braguta, P.Buividovich, T.Kalaydzhyan, S.Kusnetsov and M.Polikarpov, PoS Lattice 2010, 190 (2010); arXiv:1011,3795.

[19] M.D'Elia, S.Mukherjiee and F.Sanfilippo, Phys. Rev. D **82**, 051501 (2010); arXiv:1005.5365.

[20] M.D'Elia and F.Negro, Phys. Rev. D **83**, 114028 (2011); arXiv:1103.2080; M.D'Elia, arXiv:1209.0374.

[21] E.-M.Ilgenfritz et al., arXiv:1203.3360.

[22] G.S.Bali, F.Bruckmann, G.Enrödi, Z.Fodor, S.D.Katz and A.Schäfer, Phys. Rev. D **86**,071502 (2012); arXiv:1206.4205.

[23] Yu.A.Simonov, Phys. At. Nucl. **60**, 2069 (1997).

[24] Yu.A.Simonov, Phys. At. Nucl. **67**, 846 (2004); hep-ph/0302090; Yu.A.Simonov, Phys. At. Nucl. **67**, 1027 (2004); hep-ph/0305281.

[25] S.M.Fedorov and Yu.A.Simonov, JETP Lett. **78**, 57 (2003); hep-ph/0306216.

[26] Yu.A.Simonov, Phys. Rev. D **65**, 094018 (2002); hep-ph/0201170.

[27] Yu.A.Simonov, Phys. At. Nucl.(in press); arXiv: 1205.0692 [hep-ph].

[28] T.B.Blum, P.Chen, N.Christ et al., Phys. Rev. D **69**, 074502 (2004); V.Gimenez, V.Lubicz, F.Mescia, V.Porretti and J.Reyes, Eur. Phys. J C **41**, 535 (2005); C.Mc Neile, hep-lat/0504006; G.ColAngelo, Nucl. Phys. Proc. Suppl. **140**, 120 (2005); M.R.Pennington, hep-ph/0207220.

[29] Yu.A.Simonov and V.I.Shevchenko, arXiv:0902.1405; Yu.A.Simonov, arXiv:1003.3608; hep-ph/0501182; Yu.A.Simonov, Nucl. Phys. B **592**, 350 (2001).

[30] A.M.Badalian, B.L.G.Bakker, and Yu.A.Simonov, Phys. Rev. D **75**, 116001 (2007).

[31] M.Gell-Mann, R.L.Oakes and B.Renner, Phys. Rev. **175**, 2195 (1968).